

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-38.

LA-UR--87-49

DE87 005118

TITLE: EVALUATED DELAYED NEUTRON PRECURSOR DATA

AUTHOR(S): T. R. England, T-2
M. C. Brady, T-2
E. D. Arthur, T-2
R. J. LaBauve, T-2
F. M. Mann, Westinghouse Hanford Engineering

SUBMITTED TO: The American Nuclear Society's Annual 1987 Annual Meeting,
Dallas, Texas, June 7-11, 1987.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

MASTER

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

EVALUATED DELAYED NEUTRON PRECURSOR DATA

by

T. R. England, M. C. Brady, E. D. Arthur, R. J. LaBauve, and F. M. Mann*

Applied Nuclear Science Group
Los Alamos National Laboratory
Los Alamos, NM 87545

Based on energetics, ~ 270 fission products should be delayed neutron precursors. Of these, ~ 85 have measured and evaluated emission probabilities (Pn's) and ~ 34 have measured spectra data; the latter account for 80% or more of the total $\bar{\nu}_d$. Although the measurements are probably adequate for most reactor applications,^{1,2} complete data sets are needed for ENDF/B and for fission product yield evaluations. This paper briefly describes the use of systematics and nuclear models used to produce a complete set of data for 272 precursors and presents aggregate summation calculations needed to produce the conventional six-time-group approximations.³

Unmeasured Pn values are calculated using a fit of the 85 evaluated measurements⁴ to the Herrmann-Kratz systematic equation:⁵

$$P_n = a[(Q_\beta - S(n)/Q_\beta - K]^b \quad (1)$$

where

K = 0 for even-even precursors

= $13/A^{1/2}$ odd precursors

= $26/A^{1/2}$ odd-odd precursors

Q_β = total beta decay energy of the precursor

S(n) = neutron separation energy of daughter

a, b = fitting parameters based on measured spectra.

For unmeasured spectra we have used a statistical model code, BETA,⁶ and a simple evaporation model:

$$N(E) = CE \exp(-E/T) \quad (2)$$

where

$$[Q_\beta - S(n)] \approx a T^2 \quad (3)$$

a = fitting parameter $\approx (2/3)A$

A = mass number of precursor

* Westinghouse Hanford Engineering Development Laboratory, Richland, WA 99352

This simple model has the virtue of having a general shape similar to most measured precursor spectra and especially so in the energy range where most neutrons are emitted. It is used for cases in which no spectra have been measured. The BETA model code requires, in addition to mass values, the energy levels, spins, and parities of the precursor, daughter, and granddaughter; such detail is usually incomplete, so calculations must use added level values from systematics. The BETA code results are used only to augment 30 of the 34 measured spectra; here it is possible to normalize the calculations to a small range of the measurements.⁷

Figure 1 shows a typical measured spectrum⁸ and its comparison with the two models. The measured spectrum extends to ~ 2.5 MeV; however, for ^{94}Rb , there are likely neutrons emitted up to 3.5 MeV and more neutrons are probable at energies below ~ 100 keV than indicated by this measurement or by the models. The actual evaluated spectrum used for ^{94}Rb is an augmentation of this measurement using adjusted BETA model values above 2.5 MeV and recent low energy measurements⁹ below 0.2 MeV.

All measured precursor spectra were similarly examined and augmented, as needed, and the data file was extended, as indicated, to include Pn values and spectra for 272 precursors. These data, as well as other decay parameters, were then used in summation calculations for several fuels for both the total \bar{v}_d and spectra. Figure 2 illustrates a temporal comparison for the delayed neutrons vs time for three fuels. These results have been approximated into six groups for the equilibrium \bar{v}_d and a companion paper describes the six-group approximation for the corresponding spectra.^{3,10}

REFERENCES

1. G. Rudstam, "Six-Group Representations of the Energy Spectra of Delayed Neutrons from Fission," Nucl. Sci. Eng. 80, 238 (1982).
2. T. R. England, W. B. Wilson, R. E. Schenter, and F. M. Mann, "Aggregate Delayed Neutron Intensities and Spectra Using Augmented ENDF/B-V Precursor Data," Nucl. Sci. Eng. 85, 139 (1983).
3. M. C. Brady, T. R. England, and W. B. Wilson, "Few-Group Analysis of Current Delayed-Neutron Data," Trans. Am. Nucl. Soc. 53, p. 469 (1986).

4. F. M. Mann, M. Schreiber, R. E. Schenter, and T. R. England, "Evaluation of Delayed-Neutron Emission Probabilities," Nucl. Sci. Eng. 87, 418 (1984); updated by F. M. Mann in "The 1986 Evaluation of Delayed Emission Probabilities," presented at the Specialists' Meeting on Delayed Neutrons, Birmingham, England, September 15-19, 1986.
5. K. -L. Kratz and G. Herrmann, "Systematics of Neutron Emission Probabilities from Delayed Neutron Precursors," Z. Physik 263, 435 (1973).
6. F. M. Mann, C. Dunn, and R. E. Schenter, "Beta Decay Properties Using a Statistical Model," Phys. Rev. C 25,1, 524 (1982).
7. T. R. England, M. C. Brady, E. D. Arthur, R. J. LaBauve, and F. M. Mann, "Status of Evaluated Precursor and Aggregate Spectrum," presented at the Specialists' Meeting on Delayed Neutrons, Birmingham, England, September 15-19, 1986, Los Alamos National Laboratory informal document LA-UR 86-2693.
8. K. -L. Kratz, "Review of Delayed Neutron Energy Spectra," Proc. Consultants' Meet. on Delayed Neutron Properties, Vienna, Austria, March 26-30, 1979 [Int. Atomic Energy Agency report INDC (NDS-107/G + Special (1979))]. Also see Annual Progress Reports from the Institute for Chemistry, University of Mainz, Mainz, West Germany, 1973-1984; and personal communication K. -L. Kratz, October 1983).
9. R. C. Greenwood and A. J. Caffrey, "Delayed-Neutron Energy Spectra of ^{93}Rb and ^{143}Cs ," Nucl. Sci. Eng. 91, 305 (1985).
10. M. C. Brady and T. R. England, "Few-Group Representation of the Energy Spectra of Delayed Neutrons," submitted to the American Nuclear Society's 1987 Annual Meeting, Dallas, Texas, June 7-11, 1987.




Fig. 1 Delayed Neutron Spectra for Nuclide ^{94}Rb

Los Alamos




Fig. 2 Comparison of Total ν_d /s Following a Fission Pulse

Los Alamos